



Published in final edited form as:

*Br J Psychiatry*. 2023 July ; 223(1): 273–279. doi:10.1192/bjp.2022.185.

## Childhood Trauma, Perceived Stress, and Anhedonia in Individuals at Clinical High Risk for Psychosis

Kathleen J. O'Brien<sup>1</sup>, Arielle Ered<sup>1</sup>, Stephanie A. Korenic<sup>1</sup>, Thomas M. Olino<sup>1</sup>, Jason Schiffman<sup>2,3</sup>, Vijay A. Mittal<sup>4</sup>, Lauren M. Ellman<sup>1,a</sup>

<sup>1</sup>Temple University, Department of Psychology and Neuroscience

<sup>2</sup>University of Maryland Baltimore County, Department of Psychology

<sup>3</sup>University of California Irvine, Department of Psychology

<sup>4</sup>Northwestern University, Department of Psychology

### Abstract

**Background:** Evidence suggests that both childhood trauma and perceived stress are risk factors for the development of psychosis, as well as negative symptoms, such as anhedonia. Previous findings link increases in perceived stress to anhedonia in individuals at clinical-high risk for psychosis (CHR) and depression; however, the role of childhood trauma in this relationship has not yet been explored, despite consistent evidence that it is associated with sensitization to later stress.

**Aims:** This study examined whether perceived stress mediates the relationship between childhood trauma and anhedonia in a group of youth at CHR as well as in controls (groups with depression and no diagnosed mental health concerns).

**Methods:** The current study used multi-group mediation to examine the indirect effects of childhood trauma on anhedonia via perceived stress for CHR ( $n = 117$ ), depression ( $n = 284$ ), and non-psychiatric controls ( $n = 124$ ).

**Results:** Perceived stress mediated the relationship between childhood trauma and consummatory anhedonia regardless of group status. Perceived stress mediated the relationship between childhood trauma and anticipatory anhedonia for the CHR and depression groups, but not for non-psychiatric controls. Further, groups differed in the magnitude of this relationship, with the effects trending toward stronger for those in the CHR group.

<sup>a</sup>Corresponding Author: 1701 N 13<sup>th</sup> St Philadelphia PA 19122, ellman@temple.edu.

Author Contribution

**Kathleen J. O'Brien:** Conceptualization, Methodology, Formal Analysis, Investigation, Writing Original Draft, Review and Editing  
**Arielle Ered:** Conceptualization, Methodology, Review and Editing **Stephanie A. Korenic:** Conceptualization, Methodology, Review and Editing **Thomas M. Olino:** Methodology, Statistical Consultant, Review and Editing **Jason Schiffman:** Conceptualization, Methodology, Investigation, Review & Editing, Funding acquisition **Vijay A. Mittal:** Conceptualization, Methodology, Investigation, Review & Editing, Funding acquisition **Lauren M. Ellman:** Conceptualization, Methodology, Investigation, Review & Editing, Funding acquisition, Supervision.

Declaration of Interest

The authors have declared that there are no conflicts of interest in relation to this study.

**Conclusions:** Our findings suggest a potential transdiagnostic pathway through which childhood trauma contributes to anhedonia across serious mental illness.

---

## Introduction

Evidence suggests that childhood trauma increases risk for the development of psychosis and is associated with symptom severity across the psychosis spectrum (1-3). Childhood trauma is also associated with earlier age of onset and higher number of hospitalizations in individuals with schizophrenia (4), as well as a higher degree of positive symptoms (5,6). Importantly, evidence suggests that childhood trauma predates the onset of psychotic symptoms, suggesting a causal relationship (1). In addition to positive symptoms, childhood trauma is also associated with negative symptoms, such as anhedonia (7). It is particularly important to understand the development of negative symptoms, as they often present prior to the onset of positive symptoms (8,9) and are typically associated with worse real-world outcomes, such as deficits in social functioning (i.e., level of social contact and ability to maintain interpersonal relationships) and role functioning (i.e., level of functioning at school or work; (10). These findings indicate that childhood trauma may contribute to psychosis-risk, as well as negative symptoms specifically, and thus represents a target for early intervention.

Anhedonia has traditionally been defined as an inability to experience pleasure and can be further divided into consummatory (i.e. in-the-moment hedonic capacity) and anticipatory (i.e., ability to anticipate future enjoyment) anhedonia. Among individuals with frank psychosis, those who have experienced childhood trauma or early adversity were more likely to report symptoms of anhedonia (11). A similar relationship between childhood trauma and anhedonia has been found in those experiencing other forms of psychopathology, such as depressive disorders (12,13), as well as in non-psychiatric controls (12). However, to our knowledge, the relationship between childhood trauma and anhedonia has yet to be examined in individuals at CHR, as prior research on this population has typically focused on the relationship between childhood trauma and positive symptoms. Further, it remains unclear whether childhood trauma is differentially associated with consummatory or anticipatory anhedonia.

One potential explanation for the link between childhood trauma and psychosis is an increase in stress sensitivity, which can occur when early exposure to stress or adversity, such as childhood trauma, increases sensitivity or reactivity to stressful events later in life, as well as increasing the likelihood that daily events will be perceived as stressful (14). This impaired stress tolerance may contribute to higher levels of negative emotions and psychotic experiences in reaction to daily life stress, which constitutes a vulnerability for developing frank psychosis (15). Additionally, perceived uncontrollability of stressful events has been linked to both consummatory and anticipatory anhedonia among CHR youth (16). Using structural equation modeling, Pelletier-Baldelli and colleagues (17) found that perceived stress directly influences both consummatory and anticipatory anhedonia in a sample enriched for psychosis (i.e., scoring above cut-offs on psychosis-risk screening questionnaires), which in turn predicts deficits in social functioning. Therefore, increases in perceived stress may be a contributing factor in the development of anhedonia.

The role of childhood trauma in the perceived stress-anhedonia relationship has not yet been explored, despite consistent evidence that childhood trauma is associated with sensitization to later stress and anhedonia (14,18,19). Therefore, the current study aimed to determine whether perceived stress mediated the relationship between childhood trauma and both forms of anhedonia (anticipatory and consummatory) for CHR individuals. We hypothesized that childhood trauma would be associated with anhedonia via perceived stress. Further, although we know anhedonia is characteristic of psychosis, it also is a hallmark feature of depression, a highly comorbid disorder in CHR individuals. Understanding distinct and overlapping risk trajectories will help to pinpoint potential treatment targets. Therefore, we examined whether the same mediating relationship is present for individuals with depressive disorders, as well as community controls, to assess specificity.

## Methods

### Participants

Study participants were young adults from three large, racially, ethnically, and socioeconomically diverse catchment areas: Philadelphia County, PA, Cook County, IL, and Baltimore County, MD. Participants ( $n = 5944$ ) were recruited from local communities and universities through flyers and online sources (e.g., social media, student courses, Craigslist) to complete a baseline assessment of self-report questionnaires online using Qualtrics (Provo, Utah). A subset of these participants ( $n = 807$ ) was invited to complete in-person semi-structured interviews based on being above the pre-determined cut-off scores on two psychosis-risk questionnaires ( 8 positive symptom items on the Prodromal Questionnaire (20) or 2 endorsements of “somewhat” or “definitely agree” on the PRIME Screen (21) or randomly selected from a pool of participants below both cut-off scores (22). Starting in May 2020, interviews were conducted remotely via Zoom with cameras required to be on due to the COVID-19 pandemic ( $n = 244$ ). There were no exclusion criteria for the study beyond being fluent in English and being within the age range of 16–30, which is based on known risk periods for psychosis.

### Ethics Statement

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures were approved by the Institutional Review Boards of Temple University (Approval No. 13359), Northwestern University (Approval No. STU00205348), and University of Maryland Baltimore Country (Approval No. YS17JS20227). Further, written informed consent was obtained from all participants.

### Measures

Childhood trauma was assessed using the Childhood Trauma Questionnaire Short-Form (CTQ; (23). This is a self-report inventory assessing five types of childhood maltreatment (Emotional Abuse, Physical Abuse, Sexual Abuse, Emotional Neglect, and Physical Neglect) occurring before the age of 16. The CTQ has shown validity in both clinical and community samples for ages 12 and up, as well as convergence with the Childhood

Trauma Interview (24,25). In the current study, two of the three sites removed items that could lead to reportable situations (i.e., physical and sexual abuse) due to the online nature of the questionnaire and potential inability to contact participants. Therefore, the remaining 12 items were summed to create a modified total score ( $\alpha=.87$ ) which assesses emotional abuse, emotional neglect, and physical neglect. A subset of the sample ( $n = 255$ ) was given the full CTQ-Short Form, allowing us to determine a Pearson correlation of 0.89 between the full questionnaire and shortened version.

Perceived stress was measured by the Perceived Stress Scale (PSS; (26), which is a 14-item self-report scale assessing the extent to which daily life events in the past month are viewed as stressful and uncontrollable. The PSS has shown good reliability and validity (27-29) and has successfully discriminated between psychosis populations and controls (30-32). The total PSS score ( $\alpha=.89$ ) was used.

Anhedonia was assessed with the Temporal Experience of Pleasure Scale (TEPS; 46). The TEPS is a measure of an individual's disposition to experience pleasure and has subscales for both anticipatory pleasure (e.g., pleasure associated with expectation of reward; TEPS-ANT) and consummatory pleasure (e.g., pleasure derived while engaged in an activity; TEPS-CON). The TEPS has exhibited good internal consistency (Cronbach's alphas of 0.71-.80; (33-35), high test-retest reliability (33,35) and strong construct and discriminant validity (33,34,36). In the current study, the TEPS was used as a measure of anhedonia where lower scores indicate higher anhedonia. TEPS-ANT ( $\alpha = .76$ ) and TEPS-CON ( $\alpha = .72$ ) were our variables of interest.

CHR status was determined by the Structured Interview for Psychosis-Risk Syndromes (SIPS; 50). The SIPS is a semi-structured interview which assesses psychosis-risk syndromes and has exhibited predictive validity of conversion to psychosis (38), as well as specificity and inter-rater reliability (39-41). In the current study, all SIPS interviews were administered by individuals who had undergone extensive training led by a SIPS-certified trainer and met reliability standards ( $ICC > .80$  on positive symptom ratings). Additionally, any CHR status was verified during a weekly cross-site consensus meeting led by a SIPS-certified trainer. Participants were considered CHR ( $n = 117$ ) if they met criteria for at least one of three psychosis-risk syndromes: attenuated positive symptom syndrome (APSS;  $n = 116$ ), brief intermittent psychotic syndrome (BIPS;  $n = 0$ ), and genetic risk and functional decline (GRD;  $n = 2$ ). To address the issue of comorbidity, participants were excluded from the CHR group if they also met for a current depressive disorder ( $n = 24$ ).

The Structured Clinical Interview for DSM-5, Research Version (SCID; (42) has been termed the "gold standard" for determining clinical diagnoses. In the current study, all SCID diagnoses were confirmed in weekly meetings with a clinical supervisor. The Mood Disorders module was used to determine lifetime history of depression, including Major Depressive Disorder, Persistent Depressive Disorder, and Other Specified Depressive Disorder. Presence vs. absence of lifetime depressive disorder was the variable of interest to determine group status ( $n = 286$ ). Individuals were considered community controls ( $n = 124$ ) if they did not meet criteria for any current or past SCID diagnosis. Comorbid diagnoses for the clinical groups (CHR and depression) can be found in Table 1.

## Data Analysis

First, the dependent variables (TEPS-ANT and TEPS-CON) were examined for normality statistically by examining skewness and kurtosis values and by visually inspecting the data. Next, bivariate analyses using Pearson's correlations were conducted to determine whether there were significant relationships between the main independent variable (childhood trauma) and the potential mediator (perceived stress) and the dependent variables (TEPS-ANT and TEPS-CON), as well as if the potential mediator (perceived stress) was associated with the dependent variables (TEPS-ANT, TEPS-CON; (43). Additionally, age and sex assigned at birth (termed sex, herein) were tested as potential covariates by determining if they were associated with the independent and dependent variables (childhood trauma and anhedonia, respectively).

Statistical analyses were conducted in R (44) using version 3.6.2 of RStudio (45). The lavaan package (46) was used for analyses. Specifically, a multi-group mediation analysis for community controls, CHR, and depression groups was run so that group differences in direct and indirect effects could be examined (47). An initial model was estimated that permitted regression coefficients of all three paths to differ between groups (i.e., freely estimated). Next, a series of models were estimated which introduced constraints on individual paths by holding the regression coefficients of the path equal across groups. Each constrained model was compared to the initial freely estimated model. If the models did not significantly differ in model fit, this indicated that there were no significant differences between groups and the constrained path is preferable to the freely estimated path. Mediation effects were tested using a bootstrap estimation approach with 5,000 samples. Significant mediation was determined by the 95% confidence interval not including zero (48).

Four sensitivity analyses were run to assess consistency of results after accounting for various changes to the model parameters. First, for models in which group was a significant moderator, a multi-group mediation model was run in which all CHR individuals with past diagnoses of depressive disorders ( $n = 65$ ) were removed from the CHR group. Next, multi-group models were run with CHR individuals who also met criteria for a current depressive disorder ( $n = 24$ ) included in the CHR group. These were not run for models which were pooled across groups. Third, models were run using only the subset of individuals who completed the full CTQ-short form (i.e., including incidences of physical and sexual abuse;  $n = 152$ ). Lastly, due to the potential impact of the COVID-19 pandemic on stress and depressive symptomology, models were run with the subset of individuals assessed prior to the start of the pandemic ( $n = 359$ ).

## Results

Participant demographics and descriptive characteristics can be seen in Table 1. Compared to the depression group, the CHR group had a higher rate of Social Anxiety Disorder ( $\chi^2(1) = 6.68, p < .01$ ), but there were no other significant differences in comorbid diagnoses between the clinical groups.

All study variables were significantly correlated with each other (see Table 2). Age was correlated with TEPS-ANT ( $r = -0.08, p < 0.05$ ) and CTQ ( $r = 0.11, p < 0.01$ ) such that older age was associated with greater levels of anticipatory anhedonia and more instances of childhood trauma. Sex was correlated with TEPS-CON ( $r = 0.08, p < 0.05$ ) and PSS ( $r = 0.09, p < 0.05$ ). Therefore, to take a conservative approach, both age and sex were included as covariates in subsequent models.

First, we examined the indirect effect of childhood trauma to TEPS-ANT through perceived stress in a freely estimated model (i.e., all three paths were allowed to vary across groups) that demonstrated good model fit,  $\chi^2(21) = 138.06, p < 0.001$ ; RMSEA  $< 0.001$ , 90% CI [0.00, 0.07]; CFI = 1.00; SRMR  $< 0.001$ . We further examined the specific paths by introducing constraints on individual paths and testing for overall differences in model fit. Compared to the freely estimated model, a model with the CTQ – PSS path constrained and a model with the CTQ – TEPS-ANT path constrained did not show significantly worse fit ( $\chi^2(2) = 0.62, p = .73$  and  $\chi^2(2) = 0.74, p = .69$ , respectively), suggesting no group differences in these paths. However, a model with the PSS – TEPS-ANT path constrained was a significantly worse fit to the data ( $\chi^2(2) = 9.52, p = .01$ ), suggesting differences in this relationship across groups. Therefore, in the final model, the CTQ – PSS and CTQ – TEPS-ANT paths were constrained, and the PSS – TEPS-ANT path was allowed to vary across groups ( $\chi^2(21) = 138.06, p < 0.001$ ; RMSEA = 0.00, 90% CI [0.00, 0.07]; CFI = 1.00; SRMR = 0.01). In this model (see Figure 1), perceived stress mediated the relationship between childhood trauma and anticipatory anhedonia for the CHR and depression groups, but not for community controls. There was a significant difference in indirect effect between community controls and CHR ( $b = .10, SE = .03, z = 2.91, p < 0.01$ ), but not between community controls and depression ( $b = .04, SE = .03, z = 1.72, p = 0.09$ ). The difference in indirect effect between the CHR and depression groups was nearly significant ( $b = -.06, SE = .03, t = -1.94, p = 0.05$ ), indicating the effect was stronger for CHR at the trend level. The direct paths from CTQ to PSS ( $b = .26, SE = .03, t = 7.44, p < .001$ ) and from CTQ to TEPS-ANT ( $b = -.16, SE = .04, t = -4.74, p < .001$ ) were significant for all three groups. However, the direct path from PSS to TEPS-ANT was significant for the CHR ( $b = -.42, SE = .10, t = -4.40, p < .001$ ) and depression ( $b = -.20, SE = .06, t = -3.50, p < .001$ ) groups, but not for community controls ( $b = -.03, SE = .08, t = -0.37, p = .71$ ). Further, this relationship was significantly stronger for the CHR group compared to the depression group ( $b = -.22, SE = .12, t = -2.01, p < 0.05$ ).

For the consummatory subscale, there was no significant difference in model fit between the freely estimated model and one in which all three paths are constrained across groups, indicating there is no variation in path coefficients by group. Therefore, a single mediation model was estimated for all participants ( $\chi^2(21) = 100.342, p < 0.001$ ; RMSEA = 0.00, 90% CI [0.00, 0.05]; CFI = 1.0; SRMR = .01). In this model (see Figure 2), perceived stress mediated the relationship between CTQ and TEPS-CON. Further, CTQ was associated with both PSS ( $b = .26, SE = .03, t = 7.47, p < .001$ ) and TEPS-CON ( $b = -.07, SE = .03, t = -2.41, p = .02$ ), and PSS was also associated with TEPS-CON ( $b = -.13, SE = .04, t = -3.75, p < .001$ ).

## Sensitivity Analyses

First, a follow-up analysis of the multi-group TEPS-ANT model was run in which all individuals with past diagnoses of depression were removed from the CHR group. Although the CHR group size was significantly reduced ( $n = 52$ ), the indirect effect from childhood trauma to TEPS-ANT through perceived stress was significant. Further, there was a significant difference in indirect effects between the CHR and depression groups ( $b = -.11$ ,  $SE = .04$ ,  $t = -2.846$ ,  $p < 0.01$ ), indicating that the indirect effect was stronger for the CHR group. Next, the TEPS-ANT model was run with CHR individuals who also met criteria for a current depressive disorder ( $n = 24$ ) included in the CHR group. In this model, there was a significant difference in indirect effect between community controls and CHR ( $b = .10$ ,  $SE = .03$ ,  $t = 3.08$ ,  $p < 0.01$ ) and between the CHR and depression groups ( $b = -.06$ ,  $SE = .03$ ,  $t = -2.15$ ,  $p = .03$ ), with the effect being strongest for the CHR group. These follow-up analyses were not run for TEPS-CON, as group was not a significant moderator for TEPS-CON models and the mediation model was pooled across groups. Third, both the TEPS-ANT and TEPS-CON models were run using only the subset of individuals who completed the full CTQ-short form (i.e., including incidences of physical and sexual abuse) and the indirect effect of childhood trauma on TEPS-CON through perceived stress was no longer significant, which may have been due to loss of power. Lastly, both models were run with the subset of individuals assessed prior to the start of the pandemic and all substantive conclusions remained the same. Full results of the sensitivity analyses can be seen in the supplemental materials.

## Discussion

We demonstrated that childhood trauma is associated with higher levels of consummatory and anticipatory anhedonia via the indirect effect of perceived stress for individuals at CHR and those with depression. These findings extend existing stress-anhedonia models by including childhood trauma in the model, suggesting a transdiagnostic pathway through which childhood trauma contributes to anhedonia across the psychosis-spectrum, as well as in depressive disorders. Childhood trauma was independently associated with increases in perceived stress and anhedonia, regardless of group status. In contrast, group status moderated the relationship between perceived stress and anticipatory anhedonia, with this association being strongest for CHR individuals. Our findings provide evidence that increases in perceived stress may constitute an important target for early intervention, particularly for individuals at risk for developing psychosis who have a history of trauma.

Early adversity likely contributes to the development of psychosis in part through dysregulating the hypothalamic-pituitary-adrenal (HPA) axis, which is integral in the body's response to stress (15,18). Further, it has been suggested that altered HPA functioning, and resulting increases in stress sensitivity, may contribute to anhedonia through the release of corticotropin-releasing factor (CRF), which interferes with mesolimbic functioning, thereby reducing reward motivation (19). The findings of the current study align with previous work utilizing time-lagged multilevel modeling, which found activity-related stress (i.e., one's appraisal of their capability, control, and enjoyment of the current activity) is associated with increases in anhedonia for both CHR and first-episode psychosis (16). Similarly,

perceived uncontrollability of stressful events has been linked to anhedonia, as well as blunted midbrain dopamine responses, for individuals with depression (49). The current study adds to this literature by providing evidence that increases in perceived stress may constitute a potential mechanism through which early trauma contributes to anhedonia. Further, although this mechanism appears to be transdiagnostic, our findings suggest the relationship is strongest for CHR individuals with respect to anticipatory anhedonia.

Interestingly, there were no significant group differences in any of the paths involving consummatory anhedonia. One potential explanation would be due to limitations of the TEPS, which is prospective and hypothetical in nature. While most self-report measures require thinking retrospectively to report past incidences, the TEPS requires the participant to imagine a specific situation and predict a future emotional response, which utilizes semantic rather than experiential knowledge about emotions (50). We utilized the TEPS in the current study because it is used extensively in the psychosis-risk literature and has been recommended by the National Institute of Mental Health RDoC (Research Domain Criteria) initiative. However, it has been noted that the TEPS may reflect cognitive processes involved in the appraisal of hedonic experience as opposed to actual consumption of rewards (50). Alternatively, our findings may indicate that childhood trauma leads to consummatory anhedonia via increases in perceived stress regardless of psychopathology. It has been suggested that different neural mechanisms are involved in anticipatory and consummatory anhedonia. For example, consummatory pleasure involves initial responsiveness to reward and is mediated by opioid and GABAergic connections of the mesolimbic circuit, with projections starting in the nucleus accumbens, while anticipatory pleasure relies on reward prediction (i.e., the ability to attribute incentive value, or salience to reward-predicting cues to guide decisions; (51), which is partly moderated by dopaminergic projections from the ventral tegmental area (VTA; (52). It has been suggested that CRF interferes with mesolimbic functioning by influencing dopamine release in the VTA (19). Although we did not examine these neural mechanisms directly in this study, this body of literature could potentially explain why HPA-axis dysfunction, and associated increases in perceived stress, are particularly relevant to anticipatory anhedonia. Previous findings have established that individuals with schizophrenia show intact responsiveness to reward (i.e., consummatory pleasure), but show deficits in anticipatory pleasure (50), potentially due to difficulty integrating salient information during decision making (53). However, this distinction is less clear among CHR samples, as previous research has found individuals at CHR exhibit deficits in both consummatory and anticipatory pleasure (54,55). It has been suggested that anhedonia may not be specific to CHR, but rather associated with depression and other comorbid psychopathology (56). It should also be noted that the CC group exhibited TEPS-CON scores similar to that of the depression group. This may be reflective of rising rates of depression among young adults (57), despite not meeting clinical criteria for a depressive disorder, and highlights an important finding that should be followed up in epidemiological samples. Further, more research is needed to disentangle the mechanisms underlying anhedonia subtypes across diagnoses. However, our findings provide evidence that the link between perceived stress and anticipatory anhedonia is stronger for CHR individuals than those with depression alone.



A strength of the study was utilizing a large community sample from highly diverse populations. However, it should be noted that the current study employed community resources, such as flyers and internet sources, rather than targeting treatment-seeking populations, such as clinics or hospitals. While recruiting from non-clinical populations has been shown to reduce true risk for psychosis (58), our study used validated psychosis-risk questionnaires to select participants for clinical interviews. Further some important limitations should be noted. First, due to high comorbidity between psychosis-risk and depression, it is difficult to determine specificity of underlying mechanisms. Although it significantly reduced group size, a sensitivity analysis was conducted after removing individuals with a lifetime history of depression from the CHR group. In this sensitivity test, there was a significant difference in indirect effects between groups, indicating that the indirect effect was stronger for the CHR group. Further, the same was found when individuals with current comorbid depression were included in the CHR group, which may be a more naturalistic reflection of the CHR state. A second limitation of the study is the removal of reportable items from the CTQ (i.e., physical and sexual abuse), leaving a modified total score to be calculated using items that primarily assess neglect. However, sensitivity analyses were conducted with the subset of individuals who were given the full short form-CTQ, and the results remained the same. Additionally, among those who completed the full questionnaire, the correlation between the excluded and included items was very high ( $r = .89$ ). Third, this study relied on retrospective reporting of childhood trauma. Although the CTQ is a well-validated measure of early trauma (24,25), it has been reported that only 52% of individuals who reported traumatic events in prospective studies reported having experienced trauma when asked retrospectively (59). Therefore, the prevalence of childhood trauma may be underreported in the current study. Fourth, it is possible that our findings are not specific to anhedonia but may also apply to other negative symptoms not examined in the current study, such as motivational deficits. Finally, although the occurrence of childhood trauma precedes the other variables examined in the study, cross-sectional data does not allow for establishing temporal precedence, particularly for perceived stress and anhedonia. Longitudinal or time-lagged studies may be better able to provide support for a causal relationship between perceived stress and anhedonia. However, the temporal sequence of our study variables would be difficult to parse apart, even in a longitudinal study, given that both perceived stress and anhedonia are common responses to trauma and therefore could occur simultaneously. The current study suggests they are important constructs related to experiences of childhood trauma across groups, not only in psychosis populations.

Our findings provide evidence that childhood trauma may contribute to anhedonia via increases in perceived stress. Although this mechanism is likely transdiagnostic, it appears that the association between perceived stress and anticipatory anhedonia is strongest among CHR individuals. Therefore, the early identification and treatment of high perceived stress may represent an important intervention target, particularly for individuals at risk for developing psychosis. Cognitive-behavioral interventions, such as cognitive reappraisal and coping skills training, may be particularly beneficial, as they have been shown to reduce perceived stress (60).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Funding

This work was supported by the National Institute of Mental Health (5R01MH112612-05 to JS, 5R01MH112545-05 to VAM, 5R01MH112613-05 to LME, F31MH119720 to AE).


## References

1. Read J, van Os J, Morrison AP, Ross CA. Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica*. 2005;112(5):330–50. [PubMed: 16223421]
2. Loewy RL, Corey S, Amirfathi F, Dabit S, Fulford D, Pearson R, et al. Childhood trauma and clinical high risk for psychosis. *Schizophr Res*. 2019 Mar;205:10–4. [PubMed: 29779964]
3. Ered A, Gibson LE, Maxwell SD, Cooper S, Ellman LM. Coping as a mediator of stress and psychotic-like experiences. *Eur Psychiatry*. 2017 Jun;43:9–13. [PubMed: 28365470]
4. Rosenthal A, Meyer MS, Mayo D, Tully LM, Patel P, Ashby S, et al. Contributions of childhood trauma and atypical development to increased clinical symptoms and poor functioning in recent onset psychosis. *Early Interv Psychiatry*. 2020 Dec;14(6):755–61. [PubMed: 32043308]
5. Uyan TT, Baltacioglu M, Hocaoglu C. Relationships between childhood trauma and dissociative, psychotic symptoms in patients with schizophrenia: a case-control study. *Gen Psychiatr*. 2022 Jan 27;35(1):e100659. [PubMed: 35146333]
6. Wang Z, Xue Z, Pu W, Yang B, Li L, Yi W, et al. Comparison of first-episode and chronic patients diagnosed with schizophrenia: symptoms and childhood trauma. *Early Interv Psychiatry*. 2013 Feb;7(1):23–30. [PubMed: 22947390]
7. Kirkpatrick B, Fenton WS, Carpenter WT, Marder SR. The NIMH-MATRICES Consensus Statement on Negative Symptoms. *Schizophr Bull*. 2006 Apr;32(2):214–9. [PubMed: 16481659]
8. Mason O, Startup M, Halpin S, Schall U, Conrad A, Carr V. Risk factors for transition to first episode psychosis among individuals with “at-risk mental states.” *Schizophr Res*. 2004 Dec 1;71(2–3):227–37. [PubMed: 15474894]
9. Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell’Olio M, et al. Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. *Aust N Z J Psychiatry*. 2005 Dec;39(11–12):964–71. [PubMed: 16343296]
10. Devoe DJ, Braun A, Seredynski T, Addington J. Negative Symptoms and Functioning in Youth at Risk of Psychosis: A Systematic Review and Meta-analysis. *Harv Rev Psychiatry*. 2020 Dec;28(6):341–55. [PubMed: 33156155]
11. Sweeney S, Air T, Zannettino L, Galletly C. Gender Differences in the Physical and Psychological Manifestation of Childhood Trauma and/or Adversity in People with Psychosis. *Front Psychol*. 2015;6:1768. [PubMed: 26635676]
12. Fan J, Liu W, Xia J, Li S, Gao F, Zhu J, et al. Childhood trauma is associated with elevated anhedonia and altered core reward circuitry in major depression patients and controls. *Hum Brain Mapp*. 2021 Feb 1;42(2):286–97. [PubMed: 33030766]
13. Sonmez A, Lewis C, Athreya A, Shekunov J, Croarkin P. Preliminary Evidence for Anhedonia as a Marker of Sexual Trauma in Female Adolescents. *Adolescent Health, Medicine and Therapeutics*. 2021 Jun 1;Volume 12:67–75. [PubMed: 34163277]
14. Collip D, Myin-Germeys I, Van Os J. Does the Concept of “Sensitization” Provide a Plausible Mechanism for the Putative Link Between the Environment and Schizophrenia? *Schizophrenia Bulletin*. 2008 Mar 1;34(2):220–5. [PubMed: 18203757]
15. Myin-Germeys I, van Os J. Stress-reactivity in psychosis: evidence for an affective pathway to psychosis. *Clin Psychol Rev*. 2007 May;27(4):409–24. [PubMed: 17222489]


16. Gerritsen C, Bagby RM, Sanches M, Kiang M, Maheandiran M, Pnce I, et al. Stress precedes negative symptom exacerbations in clinical high risk and early psychosis: A time-lagged experience sampling study. *Schizophrenia Research*. 2019 Aug 1;210:52–8. [PubMed: 31248749]
17. Pelletier-Baldelli A, Strauss GP, Kuhney FS, Chun C, Gupta T, Ellman LM, et al. Perceived stress influences anhedonia and social functioning in a community sample enriched for psychosis-risk. *J Psychiatr Res*. 2021 Mar;135:96–103. [PubMed: 33460840]
18. Raymond C, Marin MF, Majeur D, Lupien S. Early child adversity and psychopathology in adulthood: HPA axis and cognitive dysregulations as potential mechanisms. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018 Jul 13;85:152–60. [PubMed: 28751271]
19. Stanton CH, Holmes AJ, Chang SWC, Joormann J. From Stress to Anhedonia: Molecular Processes through Functional Circuits. *Trends Neurosci*. 2019 Jan;42(1):23–42. [PubMed: 30327143]
20. Loewy RL, Bearden CE, Johnson JK, Raine A, Cannon TD. The prodromal questionnaire (PQ): preliminary validation of a self-report screening measure for prodromal and psychotic syndromes. *Schizophr Res*. 2005 Nov 1;79(1):117–25. [PubMed: 16276559]
21. Miller T, Cicchetti D, Markovich P, McGlashan T, Woods S. The SIPS screen: a brief self-report screen to detect the schizophrenia prodrome. *Schizophrenia Research*. 2004;70:78–78.
22. Ellman LM, Schiffman J, Mittal VA. Community Psychosis Risk Screening: An Instrument Development Investigation. *J Psychiatr Brain Sci*. 2020;5:e200019. [PubMed: 32944657]
23. Bernstein DP, Fink L, Handelsman L, Foote J, Lovejoy M, Wenzel K, et al. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *Am J Psychiatry*. 1994 Aug;151(8):1132–6. [PubMed: 8037246]
24. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl*. 2003 Feb;27(2):169–90. [PubMed: 12615092]
25. Bernstein DP, Fink L. Childhood trauma questionnaire: A retrospective self-report. Psychological Corporation; 1998.
26. Cohen S, Kamarck T, Mermelstein R. A Global Measure of Perceived Stress. *Journal of Health and Social Behavior*. 1983;24(4):385–96. [PubMed: 6668417]
27. Cohen S. Psychosocial models of the role of social support in the etiology of physical disease. *Health Psychol*. 1988;7(3):269–97. [PubMed: 3289916]
28. Cohen S, Tyrrell DA, Smith AP. Negative life events, perceived stress, negative affect, and susceptibility to the common cold. *J Pers Soc Psychol*. 1993 Jan;64(1):131–40. [PubMed: 8421249]
29. Hewitt PL, Flett GL, Mosher SW. The Perceived Stress Scale: Factor structure and relation to depression symptoms in a psychiatric sample. *J Psychopathol Behav Assess*. 1992 Sep 1;14(3):247–57.
30. Horan WP, Brown SA, Blanchard JJ. Social anhedonia and schizotypy: The contribution of individual differences in affective traits, stress, and coping. *Psychiatry Research*. 2007;149(1–3):147–56. [PubMed: 17109970]
31. Mondelli V, Dazzan P, Hepgul N, Di Forti M, Aas M, D'Albenzio A, et al. Abnormal cortisol levels during the day and cortisol awakening response in first-episode psychosis: The role of stress and of antipsychotic treatment. *Schizophrenia Research*. 2010 Feb 1;116(2):234–42. [PubMed: 19751968]
32. Palmier-Claus JE, Dunn G, Lewis SW. Emotional and symptomatic reactivity to stress in individuals at ultra-high risk of developing psychosis. *Psychol Med*. 2012 May;42(5):1003–12. [PubMed: 22067414]
33. Gard DE, Gard MG, Kring AM, John OP. Anticipatory and consummatory components of the experience of pleasure: A scale development study. *Journal of Research in Personality*. 2006 Dec 1;40(6):1086–102.
34. Gard DE, Gard MG, Mehta N, Kring AM, Patrick CJ. Impact of motivational salience on affect modulated startle at early and late probe times. *International Journal of Psychophysiology*. 2007;66(3):266–70. [PubMed: 17566577]

35. Strauss GP, Wilbur RC, Warren KR, August SM, Gold JM. Anticipatory vs. Consummatory Pleasure: What is the Nature of Hedonic Deficits in Schizophrenia? *Psychiatry Res.* 2011 May 15;187(1-2):36–41. [PubMed: 21295860]
36. Favrod J, Ernst F, Giuliani F, Bonsack C. [Validation of the Temporal Experience of Pleasure Scale (TEPS) in a French-speaking environment]. *Encephale.* 2009 Jun;35(3):241–8. [PubMed: 19540410]
37. McGlashan T, Walsh BC, Woods SW. Structured Interview for Psychosis-Risk Syndromes. Yale School of Medicine; 2001.
38. Cannon TD, Cadenhead K, Cornblatt B, Woods SW, Addington J, Walker E, et al. Prediction of psychosis in youth at high clinical risk: a multisite longitudinal study in North America. *Arch Gen Psychiatry.* 2008 Jan;65(1):28–37. [PubMed: 18180426]
39. Miller TJ, McGlashan TH, Rosen JL, Somjee L, Markovich PJ, Stein K, et al. Prospective diagnosis of the initial prodrome for schizophrenia based on the Structured Interview for Prodromal Syndromes: preliminary evidence of interrater reliability and predictive validity. *Am J Psychiatry.* 2002 May;159(5):863–5. [PubMed: 11986145]
40. Miller TJ, McGlashan TH, Rosen JL, Cadenhead K, Cannon T, Ventura J, et al. Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, interrater reliability, and training to reliability. *Schizophr Bull.* 2003;29(4):703–15. [PubMed: 14989408]
41. Woods SW, Addington J, Cadenhead KS, Cannon TD, Cornblatt BA, Heinssen R, et al. Validity of the prodromal risk syndrome for first psychosis: findings from the North American Prodrome Longitudinal Study. *Schizophr Bull.* 2009 Sep;35(5):894–908. [PubMed: 19386578]
42. First MB, Williams JBW, Karg RS, Spitzer R. Structured Clinical Interview for DSM-5—Research Version (SCID-5 for DSM-5, Research Version; SCID-5-RV). Arlington, VA, American Psychiatric Association; 2015.
43. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol.* 1986 Dec;51(6):1173–82. [PubMed: 3806354]
44. R Development Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; 2019.
45. RStudio Team. RStudio: Integrated Development for R. Boston, MA: RStudio; 2020.
46. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software.* 2012 May 24;48:1–36.
47. Ryu E, Cheong J. Comparing Indirect Effects in Different Groups in Single-Group and Multi-Group Structural Equation Models. *Frontiers in Psychology.* 2017;8:747. [PubMed: 28553248]
48. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods.* 2008 Aug;40(3):879–91. [PubMed: 18697684]
49. Pizzagalli DA. Depression, Stress, and Anhedonia: Toward a Synthesis and Integrated Model. *Annu Rev Clin Psychol.* 2014 Mar 28;10:393–423. [PubMed: 24471371]
50. Strauss GP, Gold JM. A new perspective on anhedonia in schizophrenia. *Am J Psychiatry.* 2012 Apr;169(4):364–73. [PubMed: 22407079]
51. Waltz JA, Frank MJ, Robinson BM, Gold JM. Selective reinforcement learning deficits in schizophrenia support predictions from computational models of striatal-cortical dysfunction. *Biol Psychiatry.* 2007 Oct 1;62(7):756–64. [PubMed: 17300757]
52. Barch DM, Pagliaccio D, Luking K. Mechanisms Underlying Motivational Deficits in Psychopathology: Similarities and Differences in Depression and Schizophrenia. *Curr Top Behav Neurosci.* 2016;27:411–49. [PubMed: 26026289]
53. Heerey EA, Bell-Warren KR, Gold JM. Decision-making impairments in the context of intact reward sensitivity in schizophrenia. *Biological Psychiatry.* 2008;64(1):62–9. [PubMed: 18377874]
54. Schlosser DA, Fisher M, Gard D, Fulford D, Loewy RL, Vinogradov S. Motivational deficits in individuals at-risk for psychosis and across the course of schizophrenia. *Schizophr Res.* 2014 Sep;158(0):52–7. [PubMed: 25008792]

55. Strauss GP, Ruiz I, Visser KH, Crespo LP, Dickinson EK. Diminished Hedonic response in neuroleptic-free youth at ultra high-risk for psychosis. *Schizophr Res Cogn*. 2017 Dec 23;12:1–7. [PubMed: 29928593]
56. Akouri-Shan L, Schiffman J, Millman ZB, Demro C, Fitzgerald J, Rakhshan Rouhakhtar PJ, et al. Relations Among Anhedonia, Reinforcement Learning, and Global Functioning in Help-seeking Youth. *Schizophrenia Bulletin*. 2021 Nov;47(6):1534–43. [PubMed: 34240217]
57. Goodwin RD, Dierker LC, Wu M, Galea S, Hoven CW, Weinberger AH. Trends in U.S. Depression Prevalence From 2015 to 2020: The Widening Treatment Gap. *American Journal of Preventive Medicine*. 2022 Sep;S0749379722003336.
58. Fusar-Poli P, Schultze-Lutter F, Cappucciati M, Rutigliano G, Bonoldi I, Stahl D, et al. The Dark Side of the Moon: Meta-analytical Impact of Recruitment Strategies on Risk Enrichment in the Clinical High Risk State for Psychosis. *Schizophr Bull*. 2016 May;42(3):732–43. [PubMed: 26591006]
59. Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement Between Prospective and Retrospective Measures of Childhood Maltreatment: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2019 Jun 1;76(6):584–93. [PubMed: 30892562]
60. Yusuf M, Nicoloro-SantaBarbara J, Grey NE, Moyer A, Lobel M. Meta-analytic evaluation of stress reduction interventions for undergraduate and graduate students. *International Journal of Stress Management*. 2019 May;26(2):132–45.



**Figure 1.** Multi-group structural equation model predicting anticipatory anhedonia for community controls, CHR, and depression groups. Significant indirect effect was determined by the 95% confidence interval not including zero.  $p < .001$ \*\*\*.



**Figure 2.** Structural equation model, pooled across groups, predicting consummatory anhedonia. Significant indirect effect was determined by the 95% confidence interval not including zero.  $p < .001^{***}$ .

**Table 1.**

## Sample Characteristics by Group

	<b>Control (n = 124)</b>	<b>CHR (n = 117)</b>	<b>Depression (n = 284)</b>
Age (years) M (SD) [range]	20.38 (2.82) [16-30]	20.36 (2.35) [16-30]	21.19 (2.95) [16-30]
Sex % (n) Male	30.65% (38)	29.1% (34)	16.55% (47)
Ethnicity % (n) Hispanic	13.71% (17)	10.26% (12)	20.07% (57)
Race % (n)			
American Indian/Alaska Native	0	0.9% (1)	0
Asian	29.03% (36)	17.1% (20)	22.2% (63)
Native Hawaiian/Pacific Islander	0	0.9% (1)	0.35% (1)
Black/African American	16.12% (20)	16.2% (19)	11.6% (33)
White	46.0% (57)	56.4% (66)	54.6% (155)
More than One race	7.26% (9)	3.41% (4)	8.45% (24)
Unknown/Not reported	1.61% (2)	5.13% (6)	2.81% (8)
CTQ Total Score M (SD) [range]	20.40 (6.81) [13-47]	29.17 (10.76) [13-61]	27.24 (11.06) [13-59]
PSS Total Score M (SD) [range]	23.14 (8.08) [5-50]	32.83 (8.41) [11-51]	31.79 (8.05) [6-55]
TEPS – ANT M (SD) [range]	47.38 (7.06) [22-60]	42.93 (9.36) [13-57]	44.49 (8.22) [14-60]
TEPS- CON M (SD) [range]	38.33 (6.55) [19-48]	35.98 (6.75) [20-48]	38.14 (6.37) [14-48]
Lifetime DSM-5 Diagnoses % (n)			
Alcohol Use Disorder	-	23.1% (27)	19.7% (56)
Cannabis Use Disorder	-	19.7% (23)	19.0% (54)
Panic Disorder	-	13.7% (16)	13.0% (37)
Social Anxiety Disorder	-	50.4% (59)	36.6% (104)
Generalized Anxiety Disorder	-	41.9% (49)	39.1% (111)
Obsessive-Compulsive Disorder	-	14.5% (17)	10.6% (30)
Post-Traumatic Stress Disorder	-	17.1% (20)	20.8% (59)



**Table 2.**

## Correlations Among Study Variables

	TEPS-ANT	TEPS-CON	CTQ	PSS
TEPS-ANT	1	-	-	-
TEPS-CON	.55**	1	-	-
CTQ	-.31**	-.20**	1	-
PSS	-.23**	-.13**	.36**	1

Note.

\*\*  
 $p < .01$ 

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript